

# LIVING WELL WITH MIGRAINE: BEHAVIOR AND LIFESTYLE

## What Behavioral Factors Contribute To Migraine?



### Depression

About 40% of people with migraine also have depression.

- People with migraine are twice as likely to have considered suicide than people without migraine<sup>3</sup>
- More than 66% of people with bipolar disorder have migraine<sup>4</sup>



### Anxiety<sup>15</sup>

More than half of people with migraine will have at least one anxiety disorder during their lives



### Stress

Stress is the body's automatic response to demands or challenges.<sup>10</sup>



- 4 out of every 5 people with migraine report stress as a trigger<sup>5</sup>



### Sleep

- Sleep disorders are common among people with migraine<sup>1,12,13</sup>
- More than two thirds of people with chronic migraine also have insomnia<sup>1,13</sup>
- Behavioral treatments for insomnia can help you get better sleep, and may help reduce your migraine frequency<sup>1,12</sup>



### Consistent Daily Schedule

- Having a consistent mealtime, bedtime, and daily aerobic exercise are important for people with migraine<sup>5,18</sup>
- Staying consistent in these three areas has been associated with low attack frequency in people with migraine<sup>5</sup>



### Taking Medication

- Approximately half of people with migraine do not take their medications as recommended by their doctor<sup>2</sup>
- With preventive medications<sup>2,14</sup>
  - taken daily to reduce attack frequency
  - fitting them into your daily routine is a good strategy



- Both increases and decreases in stress are associated with migraine attacks<sup>6</sup>

## What Are Behavioral Migraine Treatment Options?

Behavioral treatments help people with migraine address lifestyle factors to improve migraine management.

Some behavioral treatments that have evidence showing they reduce migraine frequency include the following:

### Relaxation Training

Daily relaxation practice can help you achieve a physical and mental state of calm and reduce migraine attacks<sup>9,11</sup>



### Cognitive Behavioral Therapy<sup>11,17</sup>

Teaches you how to break this cycle to produce a better outcome.



### Biofeedback Training

Provides instant information so you can observe and modify your body's reaction to stress<sup>9,11</sup>



There are many ways to find a psychologist who can provide these treatments:

- Ask your doctor for a referral
- Use the American Psychological Association locator, searching for a "health" specialization (<http://locator.apa.org/>)

Follow us on Facebook:

<https://www.facebook.com/pages/American-Headache-Society/67380083618>  
<https://www.facebook.com/AHMAorg>  
<https://www.facebook.com/groups/MoveAgainstMigraine>

Follow us on Twitter:

@ahsheadache @amfmigraine @ahmaorg @headachejournal

[www.AmericanHeadacheSociety.org](http://www.AmericanHeadacheSociety.org) - American Headache Society (AHS) • [www.AmericanMigraineFoundation.org](http://www.AmericanMigraineFoundation.org) - American Migraine Foundation (AMF)

<http://ahma.memberclicks.net/> - American Headache and Migraine Association (AHMA) • [www.headachejournal.org](http://www.headachejournal.org) - Headache: The Journal of Head and Face Pain

References

1. Calhoun AH, Ford S, Finkel AG, Kahn KA, Mann JD. The prevalence and spectrum of sleep problems in women with transformed migraine. *Headache*. 2006;46:604-610.  
 2. Foley KA, Cady R, Martin V, Adelman J, Diamond M, Bell CF, ... & Hu XH. Treating early versus treating mild: timing of migraine prescription medications among patients with diagnosed migraine. *Headache*. 2005; 45(5), 538-545.  
 3. Fuller-Thomson E, Schrumm M, & Brennenstuhl S. Migraine and despair: factors associated with depression and suicidal ideation among Canadian migraineurs in a population-based study. *Dep Res Treat*. 2013.  
 4. Fomaro M, and B. Stubbs. A meta-analysis investigating the prevalence and moderators of migraines among people with bipolar disorder. *J Affect Disord*. 2015; 178: 88-97.  
 5. Kelman L. The triggers or precipitants of the acute migraine attack. *Cephalalgia*. 2007; 27(5), 394-402.  
 6. Lipton RB, Buse DC, Hall CB, Tennen H, DeFreitas TA, Borkowski TM, ... & Haut SR. Reduction in perceived stress as a migraine trigger: Testing the "let-down headache" hypothesis. *Neurology*. 2014; 82(16), 1395-1401.  
 7. Martin PR, Callan M, Reece J, MacLeod C, Kaur A, Gregg K, & Goatsby PJ. Behavioral management of the triggers of recurrent headache: A randomized controlled trial. *Behav Res Ther*. 2014; 61, 1-11.  
 8. Nicholson RA, Houle TT, Rhudy JL, Norton PJ. Psychological risk factors in headache. *Headache*. 2007 Mar 1;47(3):413-26.

7. Penzien DB, Iby MB, Smitherman TA, Rains JC, Houle TT. Well-established and empirically supported behavioral treatments for migraine. *Curr Pain Headache Rep*. 2015 Jul 1;19(7):1-7.  
 8. Preutka SJ. What turns on a migraine? A systematic review of migraine precipitating factors. *Curr Pain Headache Rep*. 2014; 18(10), 1-6.  
 9. Rains JC, Poceta JS, Headache and sleep disorders: Review and implications for headache management. *Headache*. 2005; 45(2), 592-5109.  
 10. Rains JC, Poceta JS. Headache and sleep disorders: Review and implications for headache management. *Headache*. 2006;46:1344-1363.  
 11. Sancisi E, Cevoli S, Vignatelli L, et al. Increased prevalence of sleep disorders in chronic headache: A case-control study. *Headache*. 2010;50:1444-1472.  
 12. Seng EK, Rains JA, Nicholson RA, & Lipton RB. (2015). Improving medication adherence in migraine treatment. *Curr Pain Headache Rep*. 2015; 19(6), 1-7.  
 13. Smitherman TA, Kolivas ED, Bailey JR. Panic disorder and migraine: comorbidity, mechanisms, and clinical implications. *Headache*. 2013 Jan 1;53(1):23-45.  
 14. Wöber C, Brannath W, Schmidt K, Kapitan M, Rudel E, Wessely P, & Wöber-Bingöl C. (2007). Prospective analysis of factors related to migraine attacks: the PAMINA study. *Cephalalgia*, 27(4), 304-314.  
 15. Peck KR, Smitherman TA, & Baskin SM. Traditional and alternative treatments for depression: implications for migraine management. *Headache*. 2015; 55(2), 351-355.  
 16. Woldeamanuel YW, Cowan RP. The impact of regular lifestyle behavior in migraine: a prevalence case-referent study. *J Neurol*. 2016; 263: 669-676. World Health Organization. Headache disorders. Fact sheet no.277, 2012.